

Padova, 20 novembre 2018

COSA SI “ASPETTANO” I NEONATI DAGLI OGGETTI CHE LI CIRCONDANO? Fin dalle prime ore di vita integrano tutte le informazioni sensoriali in loro possesso

È stato pubblicato sulla rivista « Current Biology» lo studio del team internazionale di ricerca della Professoressa Teresa Farroni del Dipartimento di Psicologia dello Sviluppo e della socializzazione dell'Università di Padova dal titolo “[Multisensory perception of looming and receding objects in human newborns](#)” in cui si dimostra come i neonati siano dei sofisticati elaboratori del mondo complesso e dinamico che li circonda.

Subito dopo la nascita, i neonati entrano in contatto con un mondo esterno, affollato di nuovi stimoli che li bombardano da diverse posizioni. Una delle prime sfide che i neonati devono affrontare è



quella di imparare a riconoscere e integrare quelle informazioni che, attraverso canali sensoriali diversi, comunicano loro la presenza di oggetti in movimento nello spazio che li circonda. Percepire correttamente dunque degli oggetti che si muovono in direzione del proprio corpo, così, come in allontanamento è un aspetto di fondamentale importanza per l'adattamento all'ambiente. Ma qual è il processo di sviluppo di questa abilità così necessaria alla sopravvivenza?

Nello [studio](#) del team internazionale di ricerca della **Professoressa Teresa Farroni** è stato coinvolto un gruppo di neonati a cui sono stati presentati contemporaneamente due video: in questi ultimi si vedevano degli oggetti che si avvicinavano o si allontanavano visivamente. Contemporaneamente, il neonato sentiva dei suoni che potevano simulare l'avvicinamento o l'allontanamento dell'oggetto stesso attraverso la variazione di intensità sonora. Dei due video, tuttavia, solo in uno di essi il suono simulava la direzione congruente a uno dei due stimoli visivi. Sono stati misurati i tempi di fissazione per ognuno dei due stimoli visivi ed è stato dimostrato che i tempi di fissazione visiva alla nascita sono modulati dalla congruenza multisensoriale di informazioni visive e uditive che specificano il movimento degli oggetti li circondano. L'esistenza di questa precoce competenza multisensoriale suggerisce quindi che i neonati con poche ore di vita non siano completamente naïve rispetto al contesto audiovisivo che li circonda e sottolinea inoltre che i neonati sono attivi fin da subito nell'elaborare gli stimoli ambientali.

«Le ricerche che stiamo svolgendo nel nostro laboratorio – **spiega Teresa Farroni** - dimostrano non solo che a partire dalle prime ore di vita i neonati sono in grado di integrare le informazioni

sensoriali, ma che addirittura hanno delle “aspettative” su come gli oggetti si muovano intorno a loro. I neonati quindi sono da subito dei sofisticati elaboratori del mondo complesso e dinamico che li circonda.”

Link alla ricerca: [Multisensory perception of looming and receding objects in human newborns](#)

Autori: **Giulia Orioli** (Dipartimento di Psicologia dello Sviluppo e della Socializzazione dell'Università di Padova e School of Psychology, University of Birmingham), **Andrew J. Bremner** (School of Psychology, University of Birmingham) e **Teresa Farroni** (Dipartimento di Psicologia dello Sviluppo e della Socializzazione dell'Università di Padova e School of Psychology, University of Birmingham).

Correspondence

Multisensory perception of looming and receding objects in human newborns

Giulia Orioli^{1,2,*}, Andrew J. Bremner¹, and Teresa Farroni²

When newborns leave the enclosed spatial environment of the uterus and arrive in the outside world, they are faced with a new audiovisual environment of dynamic objects, actions and events both close to themselves and further away. One particular challenge concerns matching and making sense of the visual and auditory cues specifying object motion [1–5]. Previous research shows that adults prioritise the integration of auditory and visual information indicating looming (for example [2]) and that rhesus monkeys can integrate multisensory looming, but not receding, audiovisual stimuli [4]. Despite the clear adaptive value of correctly perceiving motion towards or away from the self — for defence against and physical interaction with moving objects — such a perceptual ability would clearly be undermined if newborns were unable to correctly match the auditory and visual cues to such motion. This multisensory perceptual skill has scarcely been studied in human ontogeny. Here we report that newborns only a few hours old are sensitive to matches between changes in visual size and in auditory intensity. This early multisensory competence demonstrates that, rather than being entirely naïve to their new audiovisual environment, newborns can make sense of the multisensory cue combinations specifying motion with respect to themselves.

Newborns can discriminate between visual displays showing different motion trajectories in the depth plane, showing a visual preference for trajectories approaching their body [6]. Here, we used an intersensory matching task (for example [4,7]) to investigate whether this visual preference would be altered by concurrent auditory cues specifying the motion of a sound source towards

or away from the body. Given prior evidence of intersensory matching in human infants and macaques [4,7], we predicted that, if newborns can match changes in auditory intensity and visual size according to their motion direction, then they should look longer at the visual display matching the auditory presentation: the looming display if the sound intensity increased and the receding one if it decreased. But because newborns show a preference for looming over receding visual displays [6], the visual preference for the intersensory match might also be reduced when the auditory stimulus decreased in intensity [4].

Twenty newborns (19–90 hours old) took part in the study, comprising two sound conditions, each including two trials. In both conditions, we presented the newborns with two simultaneous, side-by-side visual displays, each presenting the same object, one looming towards and the other receding from the newborn. Each trial lasted 44 seconds and included eight repetitions of the looming/receding displays). A centrally presented sound (8000 Hz sine wave) accompanied the visual displays. The sound either increased or decreased in intensity (from 55 to 70 dB or vice versa), respectively simulating approach or withdrawal of a sound source [8] (Figure 1A and Supplemental Experimental Procedures).

In both sound conditions, we recorded the newborns' looking durations to each of the visual displays and compared these across conditions, according to whether the simulated motion direction of the visual and auditory stimuli were congruent or incongruent. We found a significant interaction between the congruency of the motion direction of the stimuli and the sound condition (Figure 1B). This interaction was explained by the newborns showing a visual preference for the congruent (looming) visual display when the sound intensity increased, but comparable looking durations at the looming and receding visual displays when it decreased. To investigate if and how the presence of a sound influenced the newborns' looking to the visual stimuli, we compared the results from each condition of the current study with those of a previous one presenting unisensory looming and receding visual displays identical to those shown here

[6]. The results of this comparison suggest the presence of a sound decreasing in intensity led to an increase in the newborns' visual attention only towards the display specifying motion in the same direction as that specified by the sound (the receding display, Figure S1), reducing the preference for visual looming that newborns typically demonstrate [6].

Overall, these results show that newborns are able to match dynamic auditory and visual stimuli based on whether they specify motion towards or away from the infant. We found that newborns showed a visual preference for the looming visual display when a congruent sound, increasing in intensity, was presented. But we cannot rely on this condition alone to draw any conclusions about newborns' early multisensory competence, as they might have preferred the looming visual display irrespective of the sound, as showed in previous research [6]. However, the previous findings, obtained with visual looming and receding stimuli, highlighted newborns preference for looming visual displays in absence of sound and therefore ruled out the possibility that the visual preference found here could be simply due to an arousal state driven by the presentation of a sound increasing in intensity [6]. If newborns' preference for the looming display was driven solely by visual information and they were not taking into account the sound, we would have expected them to show a visual preference for the looming display also in the Decreasing intensity condition. Conversely, when the sound decreased the newborns showed no significant visual preference. This demonstrates that newborns' pattern of looking durations is influenced by the motion direction of a sound source simulated by a concurrent sound and, therefore, it supports the claim that they are able to match auditory and visual displays specifying a common motion direction towards or away from themselves. The absence of a significant preference in this condition suggests that the newborns' visual preferences result from a trade-off between a preference for intersensory matches and a preference for the ethological salience of the looming visual display.

The striking early multisensory competence that newborns showed in this study demonstrates that humans




Figure 1. Experimental procedure and results.

(A) Description of the intersensory matching task. The experiment comprised two different sound conditions (sounds were always presented centrally), each including two trials. One condition presented a sound increasing in intensity, simulating a looming sound source, and the other presented a sound decreasing in intensity, simulating a receding sound source. During those auditory presentations, the newborns saw, side by side, looming and receding visual displays. The audio-visual stimuli were repeated eight times in each trial, with a 1-second interval between two subsequent stimuli and 4 seconds of blank screen before the first one, for an overall trial duration of 44 seconds. The positions of the two visual displays were counterbalanced across trials. The order of presentation of the two sound conditions and the two trials within each condition was counterbalanced across participants. (B) Looking durations (ms) towards the congruent and incongruent visual displays in each sound condition. Error bars indicate the standard error of the mean. Significant comparisons are indicated (** = $p < 0.01$). When the newborns were presented with a sound that increased in intensity (specifying the approach of a sound source), they showed a visual preference for the congruent (looming) visual display. No visual preference was observed when the sound decreased in intensity. We measured the looking time to each of the visual displays in both sound conditions. A 2×2 repeated measures ANOVA (Congruency: Congruent/Incongruent; Sound condition: Increasing intensity/Decreasing intensity) revealed a significant effect of the Interaction, $F(1, 19) = 6.652$, $p = 0.018$, $\eta^2 = 0.133$. No significant main effects of Congruency, $F(1, 19) = 3.916$, $p = 0.063$, $\eta^2 = .055$, or Sound condition, $F(1, 19) = 0.809$, $p = 0.380$, $\eta^2 = 0.008$, were observed. The absence of a significant main effect of Sound indicates that the sounds presented did not affect differently newborns' alertness and attentiveness. The newborns looked longer at the congruent visual display only in the Increasing intensity condition, $t(19) = 3.562$, $p = 0.002$, $d_z = 0.796$, and not in the Decreasing intensity condition, $t(19) = -0.653$, $p = 0.521$, $d_z = 0.146$.

arrive in the world well prepared to perceive their audiovisual environment in depth. Further studies are needed to determine precisely how newborns make sense of their new audiovisual world (which cues they match across the senses). Do they match looming visual and auditory cues on the basis of changing size/intensity, is looming perceived in common via changes in the total energy of stimulation, or are links made between changing sound intensity and other visual depth cues such as changes in surface reflection? All of these possibilities raise provocative

questions about the early origins of multisensory abilities because current influential accounts of multisensory development argue that newborns are largely limited to perceiving redundant intersensory properties [9]. Changes in size and intensity are not specified redundantly across modalities, and it is even questionable whether such a basic sense property as intensity is coded in a common format in the nervous system [10]. Irrespective of the specific cues used to perceive multisensory motion trajectories, this ability at birth represents an important point of access for making

sense of the multisensory world. The prenatal origins of this ability certainly deserve further study.

SUPPLEMENTAL INFORMATION

Supplemental Information includes experimental procedures, one table, one figure and one video, and can be found with this article online at <https://doi.org/10.1016/j.cub.2018.10.004>.

ACKNOWLEDGMENTS

We are grateful to the newborns who took part in the research and to their parents for their invaluable contribution. We are also deeply indebted to the medical and nursing staff of the Department of Paediatric Unit of the Hospital of Monfalcone for their collaboration. T.F. was supported by the Wellcome Trust (073985/Z/03/Z) and by the Pro Beneficentia Stiftung (Liechtenstein).

AUTHOR CONTRIBUTIONS

Conceptualization, G.O. and T.F.; Methodology, G.O. and T.F.; Software, G.O.; Formal Analysis, G.O. and A.B.; Investigation, G.O.; Resources, G.O.; Data Curation, G.O.; Writing – Original Draft, G.O., A.B. and T.F.; Writing – Review & Editing, G.O., A.B. and T.F.; Visualization, G.O.

REFERENCES

- Cappe, C., Thelen, A., Romei, V., Thut, G., and Murray, M.M. (2012). Looming signals reveal synergistic principles of multisensory integration. *J. Neurosci.* 32, 1171–1182.
- Cappe, C., Thut, G., Romei, V., and Murray, M.M. (2009). Selective integration of auditory-visual looming cues by humans. *Neuropsychologia* 47, 1045–1052.
- Maier, J.X., and Ghazanfar, A.A. (2007). Looming biases in monkey auditory cortex. *J. Neurosci.* 27, 4093–4100.
- Maier, J.X., Neuhoff, J.G., Logothetis, N.K., and Ghazanfar, A.A. (2004). Multisensory integration of looming signals by rhesus monkeys. *Neuron* 43, 177–181.
- Van der Stoep, N., Nijboer, T.C.W., Van der Stigchel, S., and Spence, C. (2015). Multisensory interactions in the depth plane in front and rear space: A review. *Neuropsychologia* 70, 335–349.
- Orioli, G., Filippetti, M.L., Gerbino, W., Dragovic, D., and Farroni, T. (2018). Trajectory discrimination and peripersonal space perception in newborns. *Infancy* 23, 252–267.
- Walker-Andrews, A.S., and Lennon, E.M. (1985). Auditory-visual perception of changing distance by human infants. *Child Dev.* 56, 544.
- Rosenblum, L.D., Carello, C., and Pastore, R.E. (1987). Relative effectiveness of three stimulus variables for locating a moving sound source. *Perception* 16, 175–186.
- Bahrack, L.E., and Lickliter, R. (2012). The role of intersensory redundancy in early perceptual, cognitive, and social development. In *Multisensory Development*, A. Bremner, D.J. Lewkowicz, and C. Spence (Eds.), (Oxford, England: Oxford University Press) pp. 183–205.

10. Stevens, S.S. (1957). On the psychophysical law. *Psychol. Rev.* 64, 153–181.

¹School of Psychology, University of Birmingham, Birmingham, B15 2SB, United Kingdom. ²Department of Developmental Psychology and Socialization (DPSS), University of Padua, 35131, Padova, Italy. *E-mail: g.orioli@bham.ac.uk